

Draft Genome Sequence of Non-*Vibrio parahaemolyticus* Acute Hepatopancreatic Necrosis Disease Strain KC13.17.5, Isolated from Diseased Shrimp in Vietnam

Hidehiro Kondo,^a Phan Thi Van,^b Lua T. Dang,^b Ikuo Hirono^a

Laboratory of Genome Science, Tokyo University of Marine Science & Technology, Tokyo, Japan^a; Research Institute for Aquaculture No. 1, BAC Ninh, Vietnam^b

A strain of *Vibrio* (KC13.17.5) causing acute hepatopancreatic necrosis disease (AHPND) in shrimp in northern Vietnam was isolated. Normally, AHPND is caused by *Vibrio parahaemolyticus*, but the genomic sequence of the strain indicated that it belonged to *Vibrio harveyi*. The sequence data included plasmid-like sequences and putative virulence genes.

Received 17 July 2015 Accepted 5 August 2015 Published 17 September 2015

Citation Kondo H, Van PT, Dang LT, Hirono I. 2015. Draft genome sequence of non-*Vibrio parahaemolyticus* acute hepatopancreatic necrosis disease strain KC13.17.5, isolated from diseased shrimp in Vietnam. *Genome Announc* 3(5):e00978-15. doi:10.1128/genomeA.00978-15.

Copyright © 2015 Kondo et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Ikuo Hirono, hirono@kaiyodai.ac.jp.

Cultured shrimp have a high economic value. Shrimp production had been steadily increasing until 2011, when the outbreak of a new disease, called early mortality syndrome (EMS) or acute hepatopancreatic necrosis disease (AHPND), caused severe damage to the industry (1, 2). AHPND causes mortality of as high as 100%, and the causative agent of AHPND is a strain of *Vibrio parahaemolyticus*. The sequenced genomes of strains from Thailand and Mexico have a plasmid that might contain putative virulence genes (3, 4). Virulent strains can be identified by PCR using primers based on the virulence genes (5).

Because the virulence genes were on a plasmid, virulence might be transferred not only among *V. parahaemolyticus* strains but also to different bacterial species. In Vietnam, an AHPND strain, KC13.17.5, was isolated and identified to be *Vibrio* sp., but the 16S rRNA gene sequence showed that it was not *V. parahaemolyticus*. To further characterize this strain, its genome was sequenced.

Bacterial DNA was prepared according to Sambrook and Russell (6). A mate-pair library was generated from DNA using an Illumina Nextera XT DNA sample preparation kit. The library was sequenced using Illumina MiSeq and MiSeq reagent kit version 2 (300 cycles). Sequence data were assembled with CLC Genomics Workbench version 6.5.1, and then the scaffolds were analyzed on the RAST server (7). BLASTn searches of the genome were used to find sequences homologous to known plasmid-like and putative virulence genes (8).

The genome of the strain was assembled into 42 scaffolds. Annotation by the RAST server identified 5,288 genes. By further characterization, the closest neighbor of this strain was predicted to be *Vibrio harveyi* ATCC BAA-1116. A homology search using the predicted genes also showed the higher similarity to the genes of *V. harveyi* and *Vibrio campbellii* than those of *V. parahaemolyticus*.

We previously found that contig 4 of TUMSAT_D06_S3 (63 kbp) was highly conserved among 3 AHPND strains but was not conserved in 3 non-AHPND strains (3). This sequence is likely a plasmid. In the present study, some of the scaffolds, such as scaffold 21 (29 kbp) and scaffold 27 (16 kbp), were almost identical to contig 4 of TUMSAT_D06_S3. The putative virulence genes were

identified and used to diagnose AHPND by PCR (5). Because scaffold 25 (3.8 kbp) was identical to the virulence genes, we speculate that this strain was made virulent by acquiring the plasmid.

Here, we sequenced an AHPND-causing species of *Vibrio* that was not *V. parahaemolyticus*. This strain also possesses the putative toxin genes and related plasmid-like sequences. The strain was closest to *V. harveyi*, meaning that the toxin genes could be transmitted to different *Vibrio* species. Further research on how the genes are transmitted across species is needed to prevent the spread of AHPND.

Nucleotide sequence accession numbers. The partial genome sequences of the strain have been deposited in DDBJ/EMBL/GenBank under the accession numbers BBXN01000001 to BBXN01000052. The raw data for the shotgun sequencing have been deposited in DDBJ Sequence Read Archive with accession no. DRA003689.

ACKNOWLEDGMENTS

This research was funded by the Ministry of Agriculture and Rural Development (MARD) of Vietnam under the project entitled “Studies on causative agent of AHPND in Northern Vietnam” and the Heiwa Nakajima Foundation, Japan, and supported by the Japan Science and Technology Agency/Japan International Cooperation Agency, Science and Technology Research Partnership for Sustainable Development (JST/JICA, SATREPS).

REFERENCES

1. Flegel TW. 2012. Historic emergence, impact and current status of shrimp pathogens in Asia. *J Invertebr Pathol* 110:166–173. <http://dx.doi.org/10.1016/j.jip.2012.03.004>.
2. Tran L, Nunan L, Redman RM, Mohny LL, Pantoja CR, Fitzsimmons K, Lightner DV. 2013. Determination of the infectious nature of the agent of acute hepatopancreatic necrosis syndrome affecting penaeid shrimp. *Dis Aquat Organ* 105:45–55. <http://dx.doi.org/10.3354/dao02621>.
3. Kondo H, Tinwongger S, Proespraiwong P, Mavichak R, Unajak S, Nozaki R, Hirono I. 2014. Draft genome sequences of six strains of *Vibrio parahaemolyticus* isolated from early mortality syndrome/acute hepatopancreatic necrosis disease shrimp in Thailand. *Genome Announc* 2(2):e00221-14. <http://dx.doi.org/10.1128/genomeA.00221-14>.
4. Gomez-Gil B, Soto-Rodríguez S, Lozano R, Betancourt-Lozano M. 2014. Draft genome sequence of *Vibrio parahaemolyticus* strain m0605, which

- causes severe mortalities of shrimps in Mexico. *Genome Announc* 2(2): e00055-14. <http://dx.doi.org/10.1128/genomeA.00055-14>.
5. Tinwongger S, Proespraiwong P, Thawonsuwan J, Sriwanayos P, Kongkumnerd J, Chaweepeak T, Mavichak R, Unajak S, Nozaki R, Kondo H, Hirono I. 2014. Development of PCR diagnosis for shrimp acute hepatopancreatic necrosis disease (AHPND) strain of *Vibrio parahaemolyticus*. *Fish Pathol* 49:159–164. <http://dx.doi.org/10.3147/jsfp.49.159>.
 6. Sambrook J, Russell DW. 2001. *Molecular cloning: a laboratory manual*, 3rd ed. Cold Spring Harbor Laboratory Press, New York, NY.
 7. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
 8. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:421. <http://dx.doi.org/10.1186/1471-2105-10-421>.